

Reproductive Biotechnologies for Improvement of Buffalo: The Current Status

G. N. Purohit*, G. P. Duggal¹, D. Dadarwal², Dinesh Kumar³, R. C. Yadav and S. Vyas⁴

Department of Veterinary Obstetrics and Gynaecology, Rajasthan Agricultural University, Bikaner 334001, India

ABSTRACT : Reproductive biotechnologies continue to be developed for genetic improvement of both river and swamp buffalo. Although artificial insemination using frozen semen emerged some decades back, there are still considerable limitations. The major problem appears to be the lack of efficient methods for estrus detection and timely insemination. Controlled breeding experiments in the buffalo had been limited and similar to those applied in cattle. Studies on multiple ovulation and embryo transfer are essentially a replica of those in cattle, however with inherent problems such as lower number of primordial follicles on the buffalo ovary, poor fertility and seasonality of reproduction, lower population of antral follicles at all stages of the estrous cycle, poor endocrine status and a high incidence of deep atresia in ovarian follicles, the response in terms of transferable embryo recovery has remained low with 0.51 to 3.0 per donor and pregnancy rates between 15 to 30%. *In vitro* production of buffalo embryos is a valid alternative to recovery of embryos by superovulation. This aspect received considerable attention during the past decade, however the proportion of embryos that develops to the blastocyst stage is still around 25-30% and hence the *in vitro* culture procedures need substantial improvement. Embryo cryo-preservation procedures for direct transfer post thaw need to be developed for bubaline embryos. Nuclear transfer and embryo cloning is a technique that has received attention in various species during recent years and can be of immense value in buffaloes as they have a low rate of embryo recoveries by both *in vitro* and *in vivo* procedures. Gender pre-selection, genome analysis, gene mapping and gene transfer are a few of the techniques that have been studied to a limited extent during recent years and are likely to be included in future studies on buffaloes. Very recently, reproductive biotechnologies have been applied to feral buffaloes as well, but the results obtained so far are modest. When fully exploited they can play an important role in the preservation of endangered species. (*Asian-Aust. J. Anim. Sci.* 2003. Vol 16, No. 7 :1071-1086)

Key Words : Artificial Insemination, Buffalo, Embryo Transfer, In Vitro Fertilization, Oocyte, Superovulation

INTRODUCTION

Buffalo (*Bubalus bubalis*) constitute the mainstay of the dairy industry in middle and southern Asia. Buffalo has the potential superiority over cattle under the harsh climate of tropics, so should be genetically exploited. Specialized reproductive technologies continue to provide new opportunities for consistent improvement and safety of products from economically valuable food animals and safeguarding rare species. This has been achieved by dissemination of elite stock through artificial insemination and it is expected that the livestock industry will be dominated with progeny tested bulls screened for genetic markers of economic traits. Sex-selected semen (Johnson, 1992; Windsor et al., 1993; Sood, 1997) embryo sexing (Sood, 1997; Appa Rao and Totey, 1999) and cloning (Singla, 1996; Parnpai et al., 2000) are expected to produce animals of preferred genotype and gender. Animal biotechnology has shown its impact on genetic selection,

animal health, production and efficiency, animal nutrition and reproduction, and niche markets for specialized animal products (Heap and Spencer, 1998).

The biotechniques developed so far have been used for achieving various goals in animal production such as efficient utilization of germ cell production in males and females, worldwide exchange of genetic material through use of frozen semen and embryos (Nilmann, 1991; Ranjan and Pathak, 1993; Mahmoudzadeh et al., 1996; Cruz, 1998). Laboratory production of embryos allows mass scale production of embryos for practical and scientific purposes as well as for the exploitation of oocytes from donors (Drost et al., 1983; Totey et al., 1991; Madan et al., 1994b; Misra et al., 1994; Cruz, 1998; Galli et al., 1998). Improvement of genetic progress could be hastened by integrated breeding programs based on biotechnological procedures like sexing, cloning and *in vitro* production of embryos. New products like novel proteins in milk of farm animals can be produced by gene transfer (Brink et al., 2000). Gene transfer could be helpful for pharmaceutical protein synthesis at comparatively lower costs, improving animal health and introduction of specific traits (Bremel, 1996). Estimation of breeding values, an accurate genetic progress and eradication of genetic disorders and genetically determined diseases could be made possible by genome analysis and gene mapping (Arvindakshan et al., 1997).

Despite various attempts for effective utilization of

* Corresponding Author: G. N. Purohit, Tel: +91-151-2524876,

Fax: +91-151-2250336, E-mail: gnpvog@yahoo.co.in

¹ Opp. Univ. Gate, Talwandi Sabo, Distt. Bhatinda, Punjab, India.

² Dept. of Animal Reproduction Gynaecology & Obstetrics, Punjab Agricultural University, Ludhiana, Punjab, India.

³ Village and Post office Papra via Sarai Distt. Jhunjhunu, Rajasthan, India.

⁴ National Research Centre on Camel, Jorbeer, Bikaner Rajasthan.

Received November 20, 2002; Accepted March 24, 2003

these techniques the improvement in reproductive efficiency for buffaloes has remained marginal and the techniques are yet to become commercially viable. In this paper the focus on reproductive biotechniques and their use for genetic improvement of buffaloes are limited to artificial insemination, cryopreservation of germ cells and embryos, ovum pick up from ovaries *in situ*, and *in vitro* maturation, fertilization and culture. Prospective areas of gender pre-selection, cloning, genome analysis, gene mapping and gene transfer are also addressed.

ARTIFICIAL INSEMINATION (AI)

The earliest application of biotechnology for improved reproduction and thereby speeding up the rate of genetic improvement through the male path was artificial insemination. The AI technique makes efficient use of semen (through dilution and appropriate doses) over many females, use of semen of males having superior genotype and also increasing the rate of conception through proper estrus detection, reproductive organ management and medication. In India the first buffalo calf was born through AI in 1943 (Ranjan and Pathak, 1993). Since then, many more have born and it has expanded in rural areas with acceptability by farmers. However, much is still needed to cover the entire buffalo population. After cryopreservation of semen became viable in buffaloes in 1972, the extenders, cryoprotectant levels and freezing procedures have become standardized for good fertility and AI has since been used globally.

Semen dilution

Conventional extenders like egg yolk citrate or milk used for cryopreservation of bull spermatozoa are considered sub-optimal for buffalo semen. The most suitable extender for buffalo semen is tris-yolk-glycerol with caffeine citric acid and raffinose (Dhami et al., 1994). Although Kumar et al. (1993) suggest cow milk as a good dilutor for preserving buffalo semen, Akhtar et al., 1990 considered tris to be superior over milk. Tes-tris-yolk-glycerol and mes-tris-yolk-glycerol have been found equally good to tris dilutors (Oba et al., 1994; Chachur et al., 1997). The acceptable level of glycerol is 6-7% and an equilibration time of 2-6 h is considered optimum. Likewise the widely acceptable level of egg yolk is 20%. A slow cooling procedure (0.2-0.4°C/min) is recommended during pre-freezing processing of buffalo semen (Sansone et al., 2000). The freezing of semen is usually done by suspending the semen straws in horizontal position 1-4 cm above liquid nitrogen for 10-20 min, after which they are immersed into liquid nitrogen at -196°C. A step-wise freezing procedure using a programmable freezer is reported to yield better post-thaw recoveries (Del Sorbo et al., 1995), however the

freezability of buffalo bull semen is considered to be poor in general. The poor freezability of buffalo spermatozoa compared to bull spermatozoa is considered to be due to inherently fragile plasma membrane of buffalo spermatozoa and considered to be due to the low membrane phospholipid content and its loss during freezing-thawing (Sansone et al., 2000). Additives like cysteine hydrochloride, EDTA or membrane stabilizers like chlorpromazine hydrochloride are used to minimize enzyme leakage and acrosomal damage during freezing. The buffalo bull semen differs from bull semen in respect of several constituents. Compared to bull semen it has high contents of calcium, chloride, acid soluble phosphorus, and alkaline phosphatase, and lower total electrolytes, ascorbic acid and seminal protein. These altered ratios of constituents are considered responsible for poor buffering capacity and poor preservability of buffalo bull semen (Agarwal and Tomar, 1998). Sephadex and Sephadex ion-exchange filtration is reported to improve the quality and freezability of extended, low quality buffalo bull semen (Ahmad et al., 2003). Thawing of frozen semen is done in a water bath at 40°C for 30 seconds (Dhami et al., 1994, Vale, 1997) although it can be done at the higher temperature of 50°C for 15 seconds.

Conception rates

There has been much concern about semen preservation and establishment of global gene pools and development of an effective system of AI service delivery (Cruz, 1998). Buffaloes in the Indian sub-continent comprise many breeds having the best milk production capacity (Murrah and Nili-Ravi) and fat content (Surti, Bhadavari and Jaffarabadi). This gene pool is required to be preferentially selected and propagated.

The major constraints in the effective utilization of AI in buffaloes include seasonal nature of reproduction, seasonal decline in semen quality, improper estrus expression including silent heat, long ovulation time, and prolonged post-partum estrus. Cruz (1998) felt that low efficiency of AI in water buffaloes is more related to human factors and hence the efficiency in AI can be improved by proper detection of the estrus, proper semen handling, the appropriate management and nutrition after the insemination of buffaloes, and post partum. Likewise, Vale (1997) was of the view that difficulty in buffalo AI is the application of the method in the field and not the semen technology. With the recent techniques of estrus detection by measuring vaginal electrical resistance (Markandeya et al., 1993; Gupta, 1998; Gupta and Purohit, 2001), milk and plasma progesterone assay (Abbas et al., 1981), laparoscopy (Jainudeen et al., 1983), ultrasonography (Manik et al., 1992) etc. and better management protocols, the reproductive efficiency is likely to improve in the near future. Global priorities for trait propagation in the river

buffaloes as well as swamp buffaloes need to be fixed up and cryopreserved semen be used on a larger scale. Crossbreeding between the swamp buffalo and river breeds are often aimed at producing milk in addition to meat and draught in the crossbreds, as reported in China and the Philippines (Xiao Yongzuo, 1989; Mamongan et al., 1994).

The conception rate with natural service, chilled semen and frozen semen have been reported to be more than 60, 35-60 and 25-45%, respectively (Jainudeen and Hafez, 1993c; Agarwal and Shankar, 1994). The conception rate in Surti buffaloes, after insemination with semen frozen in different dilutors varied between 37.5% to 59.1% (Dhami and Kodagali, 1990; Dhami et al., 1994). Different cooling procedures (Dhami and Sahni, 1994) or thawing procedures (El-Amrawi, 1997) can improve conception rates up to 65%. Villa and Fabbri (1993) suggested by their trials on Italian buffaloes using semen frozen in a commercial dilutor Laiciphos® (IMV) that the high variation in the conception rates (30.5 to 57.1%) is because of the strong influence of environmental factors. The low conception rates to AI in buffaloes could also be due to the small size of uterine body compared to cows, leading to inadvertant deposition of semen into one of the uterine horns (Zicarelli et al., 1997b). According to Vale (1997), a pregnancy rate higher than 50% can be considered as good after insemination with frozen thawed buffalo bull semen. Inseminating twice during the estrus at 6-8 h is considered a good strategy to improve conception rates in buffaloes (Rao and Venkataramulu, 1994).

CONTROLLED BREEDING

Controlled breeding is mainly achieved through control of estrus, ovulation and parturition. Gordon (1997) elucidated the use of management strategies and medicaments for reproductive manipulation of buffaloes. The medication included prostaglandins and progesterone. The use of PGF₂ alpha or its analogues administered intramuscularly in two doses 11 days apart in estrus control in buffaloes (Rao, 1979; Diaz et al., 1994) or in single dose (Cruz et al., 1992) or in combination with GnRH for predetermined AI (Aquino et al., 1989) have been reported. Alternatively, intravulvo-submucosal (IVSM) route has been advocated by various workers for reduction of the dose and hence the cost (Rao and Rao, 1988; Rao and Venkatramaiah, 1989; Dhaliwal and Sharma, 1990). This route is demonstrated to have a delayed but comparable response.

Intravaginal and ear implants impregnated with progesterone have been alternative means of estrus control and synchronization (Singh et al., 1994) in the water buffalo. Such implants, coupled with other treatments have also shown promise for induction of cyclicity in buffaloes during

the summer season (Lohan et al., 2000). Similarly GnRH treatments are also known to induce emergence of a new follicular wave in the buffalo (Kohram et al., 2000). Buffaloes in the Indian sub-continent are known to have distinct reproductive rythms during certain seasons of the year. These are considered to be due to higher prolactin levels. Use of anti-prolactinaemic drugs like bromocryptine, have shown promise in suppression of summer anestrus in buffaloes and so have been better management protocols. Studies on melatonin profiles have shown remarkable seasonal difference in both buffalo cows and heifers (Borghese et al., 1994). The administration of melatonin has been reported to initiate ovarian activity during post-partum anestrus in buffaloes (Hassan et al., 2000). Further studies on this aspect are suggested. Immune regulation of ovarian function has been studied recently (Ramadan et al., 2000) and suggests that ovulation is an inflammatory process of immune origin. The importance of management in improving reproduction in buffaloes has been recognized and continuous exposure of buffalo cows to a fertile bull with grazing management enhances ovarian activity and improves conception rates in buffaloes (Abdalla, 2003). Studies on the molecular biology of reproduction and immunology of reproduction in the buffalo are likely to be taken up in near future.

MULTIPLE OVULATION AND EMBRYO TRANSFER (MOET)

This technique of *in vivo* embryo production exploits the genetic potential of females to accelerate the multiplication of superior animals. With the first successful transfer of an *in vivo* produced embryo in the rabbit (Heap, 1891), this technique of reproduction enhancement has been used in many domestic species. Although work on MOET in the water buffalo started some three decades ago (Drost et al., 1983; Parnpai et al., 1985; Alexiev et al., 1988), because of some peculiarities the results achieved so far have been modest. The first two live calves from non-surgical transfer of bubaline embryos were born in 1983 in USA (Drost et al., 1983) and later in India (Misra et al., 1988). For genetic gain in traits like milk production, MOET appears to be a promising technique. Gandhi (1994) have expressed that in comparison to progeny testing the increase in expected genetic gain per year from juvenile and adult MOET schemes was 63 and 70%.

The techniques of MOET have essentially been a replicate of those used in cows and involves multiple ovulation of females by administration of gonadotropins at certain stage (luteal phase) of the oestrus cycle followed by estrus induction, *in vivo* fertilization, non-surgical recovery of embryos and transfer to suitable synchronized recipients. Poor estrus expression and poor palpable characteristics of

the ovarian structures (Madan et al., 1991), lower number of primordial follicles on the buffalo ovary (Samad and Nasser, 1979; Danell, 1987), poor fertility and seasonality of reproduction (Drost, 1991), a lower population of antral follicles at all stages of the oestrus cycle in buffalo (Kumar et al., 1997), poor endocrine status (Palta and Madan, 1984) and a high incidence of deep atresia in ovarian follicles (Ocampo et al., 1996; Palta et al., 1998) are some of the limiting factors which have resulted in moderate responses of the MOET technology in this species. Subtle differences exist between the basic anatomy of genital organs (smaller size of ovaries in buffalo) and physiology of cattle and buffalo in terms of sexual behaviour and sexual rhythms. Ovarian function appears to be central to the difference in physiology leading to poor reproduction in this species. In addition, current knowledge of basic and seasonal patterns of follicle development in the buffalo is insufficient. Efforts at understanding the process and temporal pattern of follicular dynamics in the buffalo can help improve reproduction in this species.

Superovulation

Studies on superovulation among buffaloes have been carried out both in the river (Drost et al., 1983; Madan, 1984; Karaivanov, 1986; Mehmood et al., 1989; Alexiev, 1990; Ambrose et al., 1991; Alonso et al., 1994) and swamp buffaloes (Thungtanawat et al., 1981; Nityawardana et al., 1982; Parnpai et al., 1985; Chantaraprateep et al., 1989; Jainudeen, 1989; Venitkul, 1989) and have included work in countries such as the USA, Thailand, Bulgaria, Malaysia, Pakistan, Philippines, Egypt and India. Recently some reports have come in from countries such as Brazil (Basurelli et al., 1999; Carvalho et al., 2000) and Vietnam (Uoc et al., 1992; Nguyen et al., 1997).

Most of the studies have been oriented towards developing the most appropriate treatment for induction of multiple ovulation. These trials include tests on the type, mode, dosage and time of hormone administration. Results from such trials have in general shown that the ovulatory response in the buffalo is lower and less predictable than that achieved in cattle. Globally between 1981 to 1989 the average number of embryos recovered per donor averaged 1.16 and transferable embryos per donor 0.51 (Kamonpatana, 1990). Although the number of ovulations and total viable embryos may be as high as 4.1 and 6.0, the number of transferable embryos is lower, about 2.0 per donor per flush (Misra et al., 1994; Basurelli et al., 1999; Carvalho et al., 2000). One reason for such a discrepancy is the failure of ovulated eggs to enter the oviduct (Basurelli et al., 2000).

The effect of type and dose of hormone has been a subject of interest. Studies utilizing equine gonadotropin

(Parnpai et al., 1985; Karaivanov, 1986; Vlachov et al., 1986; Alexiev et al., 1988; Haman et al., 1992) or purified FSH, either in a declining dose (Haman et al., 1992; Joshi et al., 1992) or as a single injection (Kasiraj et al., 1992) have been conducted. However, the results demonstrate marginal differences in terms of viable transferable embryo recoveries (Misra, 1993; Basurelli et al., 1999). A general consensus is that purified FSH is superior to equine chorionic gonadotropin (PMSG) as the latter is associated with a higher incidence of unovulated follicles as also observed in cattle (Purohit et al., 2000). A recent report (Singh et al., 2000) suggests that treatment with eCG in buffaloes for superovulation results in elevated plasma inhibin from fully developed follicles which continue for a long time and culminate in inhibition of FSH leading to poor ovulation in the remaining follicles.

Studies on the time of hormone administration (Karaivanov, 1986; Mehmood et al., 1989; Beg et al., 1997; Cruz, 1998) and hormone pre-treatment have improved the embryo recoveries only marginally in the water buffalo (Cruz, 1998). Commercial scale projects involving superovulation and embryo collection in riverine buffalo have been carried out in India (Misra et al., 1994). Over a five year period, the total embryo yield improved from 1.77 to 3.83 and the total viable embryo yield from 0.92 to 2.13. Recent trials in swamp buffaloes employing 250 mg of recombinant bovine somatotropin (rBST) on day 4 of a progestagen implant followed by superovulation with FSH on days 9-11 in equal divided doses have shown no effect on the total number of embryos recovered, but significantly ($p < 0.05$) increased the number of transferable embryos from 0.8 ± 0.3 to 3.0 ± 1.0 (Songsasen et al., 1999).

The effects of repeated superovulation have shown to produce no effect on the viable embryo recovery and the future fertility of donors. High ambient temperatures (Misra et al., 1994) and the presence of a dominant follicle (Taneja et al., 1995) have been shown to have a deleterious effect on embryo recovery, and so has the summer season (Matharoo and Singh, 1994a; Matharoo and Singh, 1994b).

GnRH agonists like deslorelin can block ovulation in the buffalo subsequent to superstimulation with FSH and hence GnRH agonist-LH protocol can be used to control follicular growth and ovulation in superstimulated buffalo (Carvalho et al., 2000). Such a protocol has shown promise in improving superovulatory response and embryo recoveries in cattle (D'Occhio et al., 1999) and can form the basis of future studies, like delaying the LH surge post superovulation or suppressing the dominant follicle before superovulation to increase the embryo recoveries. A recent report (Carvalho et al., 2002) however, have shown only marginal increase in the embryo recoveries using such a protocol for the buffalo.

Embryo Harvest

Embryos are generally recovered using standard non-surgical procedures similar to those applied in cattle. However, it is recommended that the uterus be flushed on day 5 or 6 (Drost, 1991) compared to day 7 in cattle, as embryo development is more rapid in the buffalo. Early morula to hatched blastocysts are recovered in the flushings when embryos are harvested on day 6 (Misra, 1993). Embryos have also been recovered using a laparoscope (Techakumphu et al., 1995). Embryos are transferred using procedures similar to that in cattle.

Pregnancy Rates

The reported pregnancy rates are very low. During early trials, they were between 9.2 to 17.9 percent (Drost et al., 1983). In recent years, pregnancy rates have improved from 17% (Misra et al., 1994) to 26.4% (Misra et al., 1999) in India and this follows similar reports elsewhere. Amongst the factors affecting pregnancy rates, the season, side of transfer or type of estrus have shown to have no effect, whereas asynchrony between the donor and recipient, size of the corpus luteum of recipients and quality of the embryos transferred have profound effects on the pregnancy rate (Misra et al., 1999). To date, increasing the viable embryo recovery is the largest challenge before MOET becomes commercially viable. Steps to improve the pregnancy rates should include prevention of luteolysis during the first week of transfer and reduction in the incidence of embryo mortality. There is a report in the Phillipines of a riverine buffalo calf being born following transfer in a swamp buffalo (Cruz et al., 1991).

LABORATORY PRODUCTION OF EMBRYOS

Classical *in vitro* production of embryos involves *in vitro* maturation (IVM), *in vitro* fertilization (IVF) and *in vitro* culture (IVC). Many studies have been conducted on IVM and IVF and the procedures for IVC need to be improved significantly so as to obtain transferable embryos. Poor recovery of oocytes per ovary is another major constraint in IVF protocols. The first *in vitro* fertilized buffalo calf was born a decade ago in 1991 (Madan et al., 1991). Many births have taken place thereafter.

Oocyte Recovery

Competent oocytes for *in vitro* maturation, fertilization and culture (IVMFC) are usually recovered from ovaries of live or slaughtered animals. For the latter, one or a combination of methods of follicular aspiration, dissection or ovarian slicing are used. Whatever the method of oocyte recovery it is generally agreed that in most studies buffalo ovaries yield only a small number of quality oocytes (Totey et al., 1992; Madan et al., 1994a; Das et al., 1996; Kumar et

al., 1997; Samad et al., 1998) varying between 0.7 (Totey et al., 1992) to 2.4 (Kumar et al., 1997). This number is quite small compared to cattle. The reasons for such a low recovery are the same as discussed in relation to superovulation previously in this paper. One alternative to improve oocyte yield could be *in vitro* culture of pre-antral follicles. Isolation of pre-antral follicles from buffalo ovaries has been reported recently (Gupta et al., 2001) and more research on this aspect is likely to be undertaken in future. The selection of oocytes for IVM usually employs the presence and appearance of a multi-layer compact cumulus mass (Chungsoongneon and Kamonpatana, 1991; Suzuki et al., 1991; Samad et al., 1998), and homogeneous cytoplasm within the oocyte (Chauhan et al., 1998a).

Oocyte recovery from live animals involves the use of transvaginal ultrasound guided oocyte recovery (TVOR) also known as ovum pick up (OPU) for *in vitro* maturation and fertilization (Boni et al., 1994; Galli et al., 1998) or a laparotomy (Techakumphu et al., 1995). Such techniques are utilized to retrieve oocytes repeatedly from high merit female animals without sacrificing them and without disturbing their natural reproductive cycle. In a previous study the average yield of oocyte complexes was 1.33 per ovary per collection, whereas when the animals were pre-treated with FSH, oocyte recovery increased to 3.0 per ovary (Boni et al., 1994). Recent reports (Boni et al., 1997; Tavares et al., 1997; Zicarelli et al., 1997a) however, describe that oocyte recovery can be increased by prior superovulation and up to 60% of the retrieved oocytes mature following IVF thereby increasing the embryo production in the buffalo. Usually a 5 MHz ultrasound probe (Boni et al., 1994, 1996; Arlotto et al., 2000b) is used for the oocyte recovery and all follicles 6 mm or above are aspirated. Since buffaloes have 2 to 3 follicular waves (Basurelli et al., 1997) and follicular waves can be synchronized with the use of GnRH (Kohram et al., 2000) such a system can efficiently be used to increase the laboratory production of embryos in the water buffalo. The day of cycle at the time of OPU, the time of OPU relative to post-partum and the presence of a dominant follicle are known to affect the oocyte recovery (Boni et al., 1996). Live births from transfer of blastocysts derived from oocytes retrieved by OPU and matured and fertilized *in vitro* have been reported (Galli et al., 1998). In a more recent study these authors recovered 4.5 oocytes per OPU session (67 sessions on 5 buffaloes) and on *in vitro* maturation and fertilization 40.1% of the oocytes cleaved, yielding a total of 48 embryos of the freezable quality (Galli et al., 2001).

In vitro maturation

Laboratory maturation of the oocytes after recovery is an essential part of the *in vitro* embryo production system.

Table 1. Culture media and supplements used for *in vitro* maturation of buffalo oocytes and the nuclear maturation rates achieved in various studies

Basic medium	Hormone supplements	Serum source	Other supplements	Nuclear maturation rates (%)	Author
TCM 199	FSH	FCS	-	86	Singh et al., 1989
TCM 199	FSH, estradiol	FCS	-	84	Chungsoongneon and Kanonpatana, 1991
TCM 199	hCG, FSH, estradiol	HS	-	41	Jainudeen et al., 1993a
TCM 199	LH, FSH, estradiol	FCS	-	80	Totey et al., 1993b
Ham's F-10	LH, FSH, estradiol	FCS	-	80	
TCM 199	FSH, estradiol	EBS	-	72	Totey et al., 1992
Ham's F-10	LH	EBS	-	77	
TCM 199	-	EBS	-	80	Madan et al., 1994a
TCM 199	FSH-P	FBS	-	78	Chauhan et al., 1996
TCM 199	-	-	buFF	67	Chauhan et al., 1997b
TCM 199	FSH-P	FBS	-	74	
TCM 199	FSH-P	FBS	-	79	Chauhan et al., 1998b
TCM 199	-	FBS	IGF-II	72	
Han's F-10	FSH, LH	EBS	-	66	Bacci et al., 1991
TCM 199	FSH, hCG, estradiol	FCS	-	58	Lu and Hsu, 1991
TCM 199	-	ECS	-	80	Samad et al., 1998
	-	EBS	-	82	
TCM 199	-	EBS	-	75	Yadav et al., 1997
	-	-	buFF	79	
TCM 199	-	-	buFF	68-71	Tajik et al., 2000
TCM 199	-	-	-	81	Samad et al., 2000
Ham's F-10	-	-	-	75	
DPBS	-	-	-	59	
TCM	FSH, LH, estradiol	-	-	68	Purohit and Sharma, 2002c
TCM	-	-	EGF, IGF-1	78	
Waymouth MB media	-	-	EGF, IGF-1	83	
DMEM	-	-	EGF, IGF-1	78	Kumar et al., 2002
Ham's F-10	-	-	EGF, IGF-1	83	

FCS-fetal calf serum; HS-human serum; FBS-fetal bovine serum; EBS-estrus buffalo serum; ECS-estrus cow serum; buFF-buffalo follicular fluid.

For recovered oocytes, in order to acquire developmental competence they should undergo nuclear and cytoplasmic maturation. Nuclear maturation assessment involves the spontaneous resumption of meiosis leading to metaphase II after 24 h of *in vitro* culture of oocytes in a suitable medium (Singh and Majumdar, 1992; Yadav et al., 1997). Assessment utilizes the staining of oocytes after fixing (acetic methanol) and giemsa's (Chauhan et al., 1996; Das et al., 1997) or aceto-orcein (Totey et al., 1992; Madan et al., 1994b; Tajik et al., 2000) staining. Alternatively the degree and extent of cumulus cell expansion can be used for such an assessment (Chauhan et al., 1997a).

Cytoplasmic maturation can also be assessed by the developmental ability following IVF. Oocytes are generally matured by culture of a group of 10-15 cumulus oocyte complexes (COC's) in 50-100 μ l droplets of IVM medium under sterile paraffin oil at 38.5-39.0°C in a 5% CO₂ in air atmosphere with 90-95% relative humidity in a CO₂ incubator (Totey et al., 1992; Madan et al., 1994b; Chauhan et al., 1996). Our laboratory has recently developed an alternative technique of maturing buffalo oocytes by

vaginal culture in the *in vivo* bovine vagina (Purohit and Sharma, 2002a, 2002b); oocytes are filled in a semen straw which is sealed and kept in a used CIDR-B, which is then implanted in the vagina of a cow. Only 42.2% of the oocytes reached metaphase-II stage after 24 h of culture when placed in vaginal culture compared to 72.3% reaching metaphase-II in the laboratory culture in a CO₂ incubator but the ease with which this technique can be used can prove meaningful for the future.

The culture media used for maturation of buffalo follicular oocytes are known to affect their fertilization and subsequent development (Totey et al., 1992; Samad et al., 2000). Tissue culture medium 199 (TCM 199) is the most widely used, although some studies have utilized Ham's F10, Waymouth's MB media, DPBS or DMEM. The various media and supplements used in buffalo oocyte IVM and the maturation rates are presented in Table 1. Studies on the comparative efficiencies of different sera added in buffalo oocyte IVM are very few and inconclusive.

Hormones are usually added to IVM media and known to yield high maturation rates. Beneficial effects of growth

factors on buffalo oocyte IVM under serum free conditions point towards the likely presence of these factors in serum and follicular fluid. Besides above supplements, the beneficial effects of addition of a thiol compound like cysteamine to the IVM medium to improve *in vitro* embryo production efficiency in the buffalo has been explained (Gasparini et al., 2000). The percentage of embryos that developed to the compact morula and blastocyst stage was significantly higher (22.6%) when 50 μmol^{-1} of cysteamine was added to the IVM medium (TCM 199 supplemented with hormones and sera) compared to 14.9% when no cysteamine was added. Work on the development of completely defined serum free media has progressed in cattle (Reiger et al., 1995; Lonergan et al., 1996) and culture of buffalo oocytes under such controlled conditions, and media could prove meaningful in future research.

***In vitro* Fertilization**

Subsequent to *in vitro* maturation, oocytes are usually incubated with pre-capacitated sperm for *in vitro* fertilization. The incubation time of oocytes with spermatozoa in a fertilization medium varies between 6 h (Madan et al., 1994a; Chauhan et al., 1997a, 1999) to 24 h (Totey et al., 1992; Samad et al., 1998) and up to 48 h (Chungsoongneon and Kamonpatana, 1991). The *in vitro* matured, cumulus-intact oocytes are preferred for *in vitro* fertilization in the buffalo (Nandi et al., 1998).

Capacitation of spermatozoa is essential before incubation of oocytes and spermatozoa in a fertilization medium. For capacitating cattle spermatozoa heparin, follicular fluid or calcium ionophore-A23183 are used. However, for the buffalo such capacitating procedures have examined only heparin at a dose rate of 10 $\mu\text{g}/\text{ml}$ (Totey et al., 1992; Chauhan et al., 1997b). Spermatozoa are either pre-incubated with heparin (Kumar et al., 1994; Pal and Dhanda, 1994) for a period of 4 to 6 h (Pavasuthipaisit et al., 1992; Madan et al., 1994a, 1994b; Chauhan et al., 1997a, 1999; Nandi et al., 1998) or heparin is added directly to the fertilization medium (Totey et al., 1992, 1996). Capacitation in buffalo sperms can be assessed by chlortetracycline fluorescence assay (CTC) or *Pisum sativum* agglutinin fluorescence assay (FITC-PSA) (Kaul et al., 2000).

Prior to *in vitro* capacitation, spermatozoa with high motility are concentrated with removal of seminal plasma. Such concentration methods either employ a 1 h swim-up (Chungsoongneon and Kamonpatana, 1991; Pavasuthipaisit et al., 1992; Madan et al., 1994a; Chauhan et al., 1997a; Nandi et al., 1998) or a density gradient method (Totey et al., 1993a, 1993b, 1996; Purohit, 2001). The motility of spermatozoa is increased by pre-treatment of spermatozoa with a motility enhancing agent such as caffeine (Totey et al., 1992; Madan et al., 1994a, 1994b; Chauhan et al.,

1997a; Nandi et al., 1998) or theophylline (Jainudeen et al., 1993b). A mixture of penicillamine, hypotaurine and epinephrine (PHE) is also a good alternative (Totey et al., 1992, 1993a, 1996; Purohit, 2001).

The generally recommended sperm concentration for IVF is 2 million sperms/ml (Totey et al., 1992), however, sperm concentration as low as 0.7 million sperms/ml (Chungsoongneon and Kamonpatana, 1991) and as high as 8-10 million sperms/ml (Nandi et al., 1998) have both been reported.

Semen from different buffalo males are known to have different fertilizing capacity *in vitro* (Suzuki et al., 1992; Totey et al., 1993a). The basic fertilization medium used for buffalo oocytes is Tyrode's modified medium (TALP) (Pavasuthipaisit et al., 1992; Totey et al., 1992, 1996) or Brackett and Oliphant (BO) medium, however more studies have been concentrated on BO medium (Totey et al., 1992; Madan et al., 1994a; Ocampo et al., 1996; Chauhan et al., 1997a, 1999; Nandi et al., 1998). Fertilization *in vitro* is carried out in fertilization drops of a fertilization medium covered with sterile oil and incubated in a CO_2 incubator at temperatures and humidity similar to those used for IVM. The successful fertilization is determined either by staining of oocytes with 1% orcein in 45% acetic acid to locate the male and female pronucleas or by observing cleavage of fertilized oocytes 2 days post insemination. A recent report (Ocampo et al., 2001) depicts the fertilization of swamp buffalo oocytes using river buffalo sperm.

***In vitro* Culture of Cleaved Embryos**

Changing needs of the developing bovine embryo generally create a 8-16 cell developmental block which is a culture induced phenomenon that is irreversible but does not result in immediate embryonic death (Wright and Bondioli, 1981). Due to a general lack of understanding as to what caused this developmental block, and a few but highly variable results obtained, led to the advent of co-culture systems (Gandolfi and Moor, 1987; Eyestone and First, 1989) in cattle. Such a co-culture system with buffalo oviductal epithelial cells (BOEC) has been widely used in the buffalo (Chungsoongneon and Kamonpatana, 1991; Totey et al., 1992; Jainudeen et al., 1993b; Madan et al., 1994b) and known to increase embryo development to the morula and blastocyst stages. However, the proportion of cleaved embryos that develop to the blastocyst stage is still around 20-35%. A recent study utilized either cumulus cell monolayer or amniotic fluid of a developing 3 day chick embryo for co-culture of 2-4 cell zygotes. The number of zygotes that reached the expanded blastocyst stage was 10% for both the culture systems (Ocampo et al., 2001). The *in vitro* culture system requires substantial improvement in the buffalo.

EMBRYO CRYOPRESERVATION

Although some researchers have successfully frozen buffalo embryos, cryopreservation of bubaline embryos has not been attempted on a large scale. Majority of the mammalian embryos are still frozen by conventional equilibrium methods involving controlled cooling rates which require the use of freezing machines (Palasz and Mapletoft, 1996). Using such procedures, cryopreservation of buffalo embryos has been reported (Misra et al., 1992; Kasiraj et al., 1993) but pregnancy rates are very low. Transfer of 4 embryos frozen in 1.0 M glycerol to 3 recipients resulted in 1 pregnancy vs. 2 pregnancies for 5 embryo frozen in 1.4 M glycerol and transferred to 5 recipients (Misra et al., 1992). When 39 embryos were frozen in 1.4 M glycerol and transferred to equal number of recipients over a 1 year period 11 pregnancies resulted and 9 live births recorded (Kasiraj et al., 1993).

Vitrification is a recently developed technique that requires no freezing equipment and involves a physical process by which a highly concentrated solution of cryoprotectant solidifies during cooling without the formation of ice crystals. Vitrification has been used successfully for cryopreservation of embryos at various developmental stages (Nilmann, 1991). With the reports of simple ethylene glycol/ficoll/sucrose media (Mahmoudzadeh et al., 1996) bovine embryos have now been preserved and can be directly transferred post thaw, similar to semen. Such procedures are likely to be incorporated in cryopreservation of bubaline embryos, and hence make the system more easy and effective. The results achieved in experiments carried out so far give reason to believe that it is possible to create banks for deep frozen buffalo embryos. These banks could be used successfully for conservation of genetic resources in buffaloes, introduction of new breeds and rapid dissemination of high producing buffalo genotypes (Alexiev, 1990).

Recently attempts to vitrify buffalo oocytes have been made, with post vitrification recovery rate of 88-89% and morphologically normal. These oocytes could be further used for IVM and IVF (Dhali et al., 2000a, 2000b).

EMBRYO TECHNOLOGIES IN FERAL BUFFALOES

Work on *in vitro* embryo production in the feral species like the African buffalo (*Syncerus caffer*) essentially involves immobilization using etorphine and ultrasound guided OPU (Arlotto et al., 2000a; Gerber et al., 2000a). Using a new guide system, an average recovery of 2.5 oocytes per aspiration has been reported and mean number of oocytes per aspiration in buffalo cows varies from 2.5-2.7 with a range of 0 to 9 (Gerber et al., 2000b). Studies using slaughter house ovaries however, report an average

recovery of 12 COC's per ovary by slicing or mincing (Kidson et al., 2000). However, the percentage of good quality oocytes developing to morula stage was low and did not differ significantly when collected from ovaries of calves (27.3%), non-pregnant (38.9%) and pregnant buffalo cows. Only 9.7% of oocytes recovered from pregnant cows reached up to blastocyst stage and none of the oocytes from calves or non-pregnant cows developed to blastocyst stage. Studies on ovaries from slaughter house also indicate that the number of oocytes recovered per ovary are highest (9.0 \pm 2.6 per ovary) from adult non-pregnant or early pregnant, African buffalo as compared to juvenile or late pregnant buffalo ovaries (Arlotto et al., 2000a). Most of the work on IVM\F\C on feral buffaloes use media developed in cattle or water buffaloes. TCM 199 is the basic medium supplemented with hormones and serum source (Smith et al., 1999; Kidson et al., 2000; Gerber et al., 2000a). Such recent work is likely to play an important role in the preservation of wild species, like the bison. However, concentrated efforts in maturation of oocytes need to be developed substantially. Reports on the superovulation and embryo transfer in the feral buffaloes like the wood bison (*Bison bison athabasca*) which was listed as an endangered species in Canada, and African cape buffalo (*Syncerus caffer*) have appeared only recently (Othen et al., 1999) with little or no success. Similar work on the cryopreservation of epididymal sperms of African buffalo have also appeared recently (Lambrechts et al., 1999).

NUCLEAS TRANSFER AND EMBRYO CLONING

The embryos obtained through cloning can be transferred to recipient animals and many animals of similar genetic make up could be obtained. The basic procedure involves the utilization of micro-manipulation and cell fusion to transfer blastomeres of multicellular embryo or somatic cell into enucleated oocytes. The nucleus of blastomere is reprogrammed in such a way that a new embryo develops (Singla, 1997; Singla et al., 1997). With the advent of using nuclei of somatic cells (Wilmut et al., 1997) cloned embryos in rabbits, mice, sheep, goats, cattle and pigs have been produced, however, there are still considerable limitations. Recently, cloning has been reported in buffaloes (Singla, 1996; Appa Rao and Totey, 1999; Kitiyanant et al., 2000; Parnpai et al., 2000; Shi et al., 2000). Buffalo fetal fibroblasts proved more efficient than granulosa cells in generating cloned swamp buffalo embryos, due to a high cleavage rate, morula and blastocyst formation (Parnpai et al., 2000). With a low rate of *in vivo* and *in vitro* procedures for embryo recovery in buffaloes, the technique of embryo cloning has immense value in the production of buffalo embryos and is likely to receive considerable attention in future.

GENDER PRESELECTION

Male and female offsprings are preferred for different purposes. Therefore gender preselection is considered advantageous as it has a significant impact on the economics of livestock production. Certain sectors of buffalo industry may benefit from a high producing male progeny, the most notable being beef stud breeders, but mostly female progeny are preferred for milk production. Possible mechanisms of altering sex include facilitating or inhibiting the transport of either X or Y chromosome bearing sperm through the reproductive tract, preferential selection of sperm at fertilization, or sex-specific death of embryo after fertilization (Rorie, 1999). Recent success using sperm cells sorted into X and Y chromosome bearing population by flow cytometry (Johnson, 1992; Windsor et al., 1993; Muster et al., 1999) may be applied to buffalo. However, cytotoxic and mutagenic effects of this technique and reduction in fertility has been reported, due to the use of UV-excitable DNA specific stain (Johnson et al., 1987; McNutt and Johnson, 1996).

The sex of an embryo can also be determined using polymerase chain reaction (PCR). Two to four cells of an embryo blastomere serve as a sample for the sexing assay. A DNA sequence that is repeated on Y chromosome is used as target for male specific DNA amplification. Since PCR is able to detect even a single copy of target DNA, repetitive DNA sequences serve to enhance the sensitivity and consequently embryo viability by minimizing the number of blastomeres required for the sexing assay (Sood, 1997).

Biopsy of morula and blastocyst cells can occur without impairing the embryonic viability. Isolation of different Y-chromosome specific DNA fragments from bovine species has been extended for the buffalo. Buffalo Y-chromosome specific repetitive DNA (BuRY. 1) has been cloned and sequenced for sexing of buffalo embryos by PCR (Appa Rao and Totey, 1999). Likewise, duplex PCR based assay using 2-4 cells from an embryo has also been reported for sexing buffalo embryos (Sood et al., 2001). Sexing could be done within 4 h and sex was assigned to all 59 embryos used in the study. PCR method of sex determination is quick, highly accurate and easy to perform. However, reports on the birth of calves from pre-sexed embryos are lacking but should occur in near future.

GENOME ANALYSIS AND GENE MAPPING

Molecular genetics has tremendously increased the understanding of the structure function and organization of genes. Genome analysis through gene mapping has progressed in farm animals especially in pigs, cattle and chicken. In buffalo some initial progress has also been made (Arvindakshan et al., 1997; Barker et al., 1997; Simonsen et

al., 1998). In this context, determination of DNA sequences and monitoring of these sequences in individual animals at a low cost are fundamental to the DNA-diagnosis in animal breeding. However, as most traits are of quantitative nature in farm animals, it is important to identify quantitative trait loci (QTL) at the genomic level for use in genomic selection and application of gene transfer. Marker assisted selection would be useful in avoiding risk of getting undesirable side effects. Microsatellite variation amongst river buffaloes from Sri Lanka and Malaysia and swamp buffaloes from Thailand, Malaysia, Philippines and Australia revealed significant genetic differentiation between the river and swamp buffaloes (Barker et al., 1997). The two breed types were described to have diverged some 15,000 years ago. Studies of 11 populations of the African buffalo (*Syncerus caffer*) using mtDNA sequence data and analysis of variation at 6 micro-satellite data revealed high levels of genetic variability. The expected and observed values of heterozygosities in populations were high and varied within a narrow range. The high level of genetic variation was evenly distributed among populations and significant differentiation was only observed at the continental level (Simonsen et al., 1998). Results of analysis of buffalo genomic library suggest that its genome is rich in microsatellite type sequences (Silva et al., 2002).

GENE TRANSFER

Gene transfer is the basis of producing transgenic animals. The transgenics carry recombinant DNA within their genome, introduced by intentional human intervention (Wall, 1996). The transferred genes consists of a functional part and a regulatory element, the promoter. Transgenic production involves one of a number of techniques : micro-injection into the pronucleus of fertilized oocytes, DNA-transfer via electroporation using retroviral vectors, incorporation of sperm that carry the gene construct into the ovum at fertilization (usually via intra-cytoplasmic sperm injection, ICSI), and transformation of fetal fibroblasts followed by nucleus transfer (Cibelli et al., 1998). To increase the number of transgenics born per pregnancy, a multiplex PCR analysis is performed on embryo biopsies to identify transgenics. Thus all transgenic embryos could be selected for transfer into synchronized recipients (Hendolin et al., 2000).

Reports on production of transgenic buffalo are currently not available. The only report on genetic analysis reports on parentage assessment by DNA fingerprinting of a IVF buffalo calf (Mattapallil and Ali, 1998). Gene transfer in buffalo would be a long way and would be possible once traits of economic significance are identified and the buffalo genome mapped to a greater extent. The trait of muscular hypertrophy or double muscling have been located on

bovine chromosome 2 (Charlier et al., 1995). Since buffalo has a great significance in production of meat (which is superior in protein content and contains lower cholesterol compared to cattle) identification of such traits can be of immense value in gene transfer. Some genes for superiorities of buffaloes need to be identified like ability to use low quality roughages due to a reduced nitrogen output in the urine (Moran, 1983) greater tick resistance (Chantalakhana and Skunmun, 1998) and superior quality of milk casein. It remains to be resolved whether genes of buffalo could be of significance by transfer to homologous or heterologous species, or transgenic buffalo could be of significance for producing pharmaceutical proteins.

CONCLUSION

Understanding the ovarian and follicular dynamics appears to be pivotal to improve reproduction in the buffalo. AI in the water buffaloes needs to be improved by improving estrus detection and fixing up priorities for a global trait propagation. For the feral buffaloes, however, cryopreservation procedures need to be developed. Techniques of embryo harvest both *in vivo* and *in vitro* and, embryo cryopreservation need substantial improvement, but, with some species peculiarities this does not appear true at least for embryo production. *In vivo* oocyte recovery from live animals and further IVM/F/C should be encouraged. Methods to prevent atresia of follicles and development of follicles would also help in improving the embryo recovery rates. Amongst the recent technologies such as embryo and sperm sexing, cloning, gene transfer and genome mapping, embryo bisection and nucleus transfer appears to have immense value, as this can substantially help in improving embryo numbers.

REFERENCES

- Abbas, S. K., C. S. Ali, H. A. Samad, M. Arsian and M. Yaqoob. 1981. Plasma progesterone profile of oestrus cycle in buffalo heifers measured by radioimmunoassay. *Pakistan Vet. J.* 1:76-78.
- Abdalla, E. B. 2003. Improving the reproductive performance of Egyptian buffalo cows by changing the management system. *Anim. Reprod. Sci.* 75:1-8.
- Agarwal, S. K. and U. Shankar. 1994. Annual report. Livestock Production Research, Indian Veterinary Research Institute, U.P. India.
- Agarwal, S. K. and O. S. Tomer. 1998. Reproductive technologies in buffalo. Indian Veterinary Research Institute, U.P. India.
- Ahmad, Z, M. Anzar, M. Shahab, N. Ahmad and SMH. Andrabi. 2003. Sephadex and Sephadex ion exchange filtration improves the quality and freezability of low grade buffalo semen ejaculates. *Theriogenology* 59:1189-1202.
- Akhtar, T., R. A. Chaudhary, I. H. Khan and T. M. Khan. 1990. Extracellular release of hyluronidase and acrosin from buffalo bull spermatozoa extended in different extenders. *Recent Adv. Buffalo Res.* 3:75-79.
- Alexiev, A., 1990. Genetic aspects of embryo transfer in buffalo breeding. *Theriogenology* 33:613.
- Alexiev, A., K. Vlahov, C. Karaivanov, D. Kacheva, O. Palikhronov, M. Petrov, W. Nikolov, A. Drogoev and P. C. Radev. 1988. Embryo transfer in buffaloes in Bulgaria. In: *Proc. 2nd World Buffalo Congress.* vol. 2, pp. 591-595.
- Alonso, J. C., El-campo and A. Gill. 1994. Multiovarulatory effects of PMSG and FSH in female riverine buffaloes, *Proc. IVth World Buffalo Congress,* 3:561-563.
- Ambrose, J. D., S. K. Singla, S. Jaiikhani, B. S. Prakash and M. L. Madan. 1991. Superovulation responses in buffaloes (*Bubalus bubalis*) to different regimes of Folltropin. *Theriogenology* 35:181.
- App Rao, K. B. C. and S. M. Totey. 1999. Cloning and sequencing of Buffalo male specific repetitive DNA: Sexing of *In vitro* developed buffalo embryos using multiplex and Nested Polymerase chain reaction. *Theriogenology* 51:785-797.
- Aquino, E. C., P. G. Duran, H. V. Venturina, O. F. Smith, F. T. Rellm and L. C. Cruz. 1989. Effect of time of GnRH administration on ovulation response and on CL function in post partum buffalo cows treated with PGF₂ α for synchronization of estrus. *Phil J. Vet. Anim. Sci.* 17:75.
- Arlotto, T., D. Gerber, D. Cooper and S. J. Terblanche. 2000a. Follicle number and oocyte recovery from ovaries of wild African buffalo (*Syncerus caffer*). 14th International Congress on Animal Reproduction, Compendium of Abstracts. Vol. 2:175 (Abstr).
- Arlotto, T., D. Gerber, S. J. Terblanche, D. Cooper and H. J. Bertschinger. 2000b. Development of morula stage embryos after transvaginal oocyte recovery in the African buffalo (*Syncerus caffer*) *Theriogenology* 53:324.
- Arvindakshan, T. V., A. M. Nainar and K. Nachinuthu. 1997. High salt method a simple and rapid procedure for isolation of genomic DNA from buffalo (*Bubalus bubalis*) white blood cells. *Indian Journal of Experimental Biology.* 35: 903-905.
- Bacci, M. L., G. Galeati, M. Matloili, K. Boni and E. Seren, 1991. *In vitro* maturation and *in vitro* fertilization of buffalo oocytes. In *Proceedings 3rd World Buffalo Congress* verma, Bulgaria, Vol. III, 599-603.
- Barker, J. S. F., S. S. Moore, D. J. S. Hetzel, D. Evans, S. G. Tan and K. Byrene. 1997. Genetic diversity of Asian water buffalo (*Bubalus bubalis*). Microsatellite variation and a comparison with protein coding loci. *Animal Genetics.* 28: 103-115.
- Basurelli, P. S., E. H. Madureira, J. A. Visintin, R. Porto Filho, N. A. T. Carvalho, G. Campanile and I. Zicarelli. 2000. Failure of oocyte entry into oviduct in superovulated buffalo. *Theriogenology* 53:491 (Abstr).
- Basurelli, P. S., R. G. Mucciolo, R. Arruda, E. H. Madureira, R. Amaral and M. E. O. Assumpcao. 1999. Embryo recovery rate in superovulated buffalo. *Theriogenology* 51:401 (Abstr).
- Basurelli, P. S., R. G. Mucciolo, J. A. Visintin, R. P. Arruda, E. H. Madureira, C. Olivera and J. R. Molero Filho. 1997.

- Ovarian follicular dynamics during estrous cycle in buffalo (*Bubalus bubalis*). *Theriogenology* 47:1531-1547.
- Beg, M. A., P. C. Sanwal and M. C. Yadav. 1997. Ovarian response and endocrine changes in buffalo superovulated at mid luteal and late luteal stage of the estrous cycle: A preliminary report. *Theriogenology* 47:423-432.
- Boni, R., S. Roviello and L. Zicarelli. 1996. Repeated ovum pick up in Italian Mediterranean buffalo cows. *Theriogenology* 46:899-909.
- Boni, R. S., R. Di Palo, V. Barbriere and L. Zicarelli. 1994. Ovum pick up in deep anestrus buffaloes. *Proc. IV World Buffalo Congress* 3:480-485.
- Boni, R. S., S. Roviello, B. Gasparrini and L. Zicarelli. 1997. Pregnancies established after transferring embryo yielded by ovum pick up and *in vitro* embryo production in Italian buffalo cows *Proc. Vth World Buffalo Congress, Caserta, Italy*. pp. 787-792.
- Borghese, A., V. L. Barile, G. M. Terzano, A. M. Pilla and A. Parmeganni. 1994. Melatonin trend during seasons in heifers and buffalo cows. In *Proceedings 4th World Buffalo Congress, Sao Paulo, Brazil*, vol. 3, pp. 528-530.
- Bremel, R. D., 1996. Potential role of transgenesis in dairy production and related topics. *Theriogenology*. 45:51-56.
- Brink, M. F., M. D. Bishop and F. R. Pieper. 2000. Developing efficient strategies for the generation of transgenic cattle which produce biopharmaceuticals in milk. *Theriogenology* 53:139-148.
- Carvalho, N. A., P. S. Basurelli, L. Zicarelli, E. H. Madureira, J. A. Vistin, M. J. D'Occhio. 2002. Control of ovulation with a GnRH agonist after superstimulation of follicular growth in buffalo: fertilization and embryo recovery. *Theriogenology* 58:1641-1650.
- Carvalho, N. A. T., P. S. Basurelli, L. Zicarelli, E. H. Madureira, J. A. Visentin and M. J. D'Occhio. 2000. Use of GnRH agonist to control ovulation subsequent to superstimulation of follicular growth in buffalo. *Theriogenology* 53:496.
- Chachur, M. G. M., E. Oba, C. I. M. Gonzales. 1997. Equilibrium time influence on motility vigour and membrane integrity of thawed buffalo bull semen using triladyl, glycine-egg yolk and tes extenders. *Proc. 5th World Buffalo Cong. Caserta, Italy*, vol. 1, pp. 846-849.
- Chantalakhana, C. and P. Skunmun. 1998. Long term breeding strategies for genetic improvement of buffaloes in developing countries. 8th World Conference on Animal Production Korea. *Proc. Symposium Series*, 1 pp. 306-316.
- Chantrapateep, P., C. Lohachit, M. Techakumphu, G. Kalayashi, P. Virakul, A. Kumavongkrit, P. Prateep and A. Limskulm. 1989. Early embryonic development in Thai swamp buffalo (*Bubalus bubalis*). *Theriogenology* 31:1131-1138.
- Charlier, C., M. D. Bishop and W. Barendse. 1995. The mh gene causing double muscling in cattle maps to bovine chromosome 2. *Manmalian Genome* 6:788-792.
- Chauhan, M. S., P. K. Katiyar, M. L. Madan, S. K. Singla and R. S. Manik. 1996. Influence of follicle stimulating hormone on *in vitro* maturation and cleavage of buffalo oocytes after *in vitro* fertilization. *Theriogenology* 45:243 (Abstr).
- Chauhan, M. S., P. K. Katiyar, S. K. Singla, R. S. Manik and M. L. Madan. 1997a. Production of buffalo calves through *in vitro* fertilization. *Indian J. Anim. Sci.* 67:306-308.
- Chauhan, M. S., P. Palta, S. K. Das, P. K. Katiyar and M. L. Madan. 1997b. Replacement of serum and hormone additives with follicular fluid in the IVM medium: effects on maturation, fertilization and subsequent development of buffalo oocytes *in vitro*. *Theriogenology* 48:461-469.
- Chauhan, M. S., S. K. Singh, P. Palta, R. S. Manik and M. L. Madan. 1999. Effect of EGF on cumulus expansion, meiotic maturation and development of buffalo oocytes *in vitro*. *Vet. Rec.* 144:266-267.
- Chauhan, M. S., S. K. Singla, P. Palta, R. S. Manik and M. L. Madan. 1998a. *In vitro* maturation and fertilization and subsequent development of buffalo (*Bubalus bubalis*) embryo: Effect of oocyte quality and type of serum. *Reprod. Fertil. Dev.* 10:173-177.
- Chauhan, M. S., S. K. Singla, P. Palta, R. S. Manik and O. S. Tomer. 1998b. IGF-II stimulation of *in vitro* maturation, *in vitro* fertilization and subsequent development of buffalo (*Bubalus bubalis*) oocytes *in vitro*. *Vet. Rec.* 142:727-728.
- Chungsoongneon, U. and M. Kamonpatana. 1991. Oocyte maturation, *in vitro* fertilization and culture system for developing preimplantation swamp buffalo embryos using frozen-thawed semen. *Buffalo J.* 7:189-98.
- Cibelli, J. B., S. L. Stice, P. J. Golneke, J. J. Kane, J. Jery, C. Blackwell, F. A. Ponce de Leon and J. M. Robl. 1998. Cloned transgenic calves produced from quiescent fetal fibroblasts. *Science*. 280:1256-1258.
- Cruz, L. C., E. J. Aquino, P. Duran, H. Venturina, P. S. Faylon, and S. K. Ranjhan. 1991. Abnormalities of CL function in post partum water buffaloes following early induction of ovulation with GnRH. *Proc. 3rd World Buffalo Congress, Verma, Bulgaria*. 3:721.
- Cruz, L. C., O. F. Smith, F. A. Adriano, H. V. Venturina, F. Lorenzo, F. Manuad, E. Ataby and P. G. Duran. 1992. Ovarian follicular dynamics and response to superovulatory treatments in water buffaloes. *Phil. J. Vet. Anim. Sci.* 18:19-32.
- Cruz, L. C. 1998. Reproductive Biotechnique in Buffaloes. 8th World Conference on Animal Production. Symposium series. pp. 286-289.
- Danell, B. 1987. Estrus behaviour, ovarian morphology and cyclical variation in follicular system and endocrine pattern in water buffalo heifers. Thesis, Swedish University of Agricultural Sciences, Uppsala, Sweden. pp. 124.
- Das, G. K., G. C. Jain, V. S. Solanki and V. N. Tripathi. 1996. Efficacy of various collection methods for oocyte retrieval in buffalo. *Theriogenology* 46:1403-1411.
- Das, S. K., M. S. Chauhan, P. Palta and O. S. Tomer. 1997. Influence of cumulus cells on *in vitro* maturation of denuded buffalo oocytes. *Vet. Rec.* 141:552-553.
- Del Sorbo, C., A. Fabbrocini, G. Fasano, M. R. Martusciello and G. Sansone. 1995. The effect of two different freezing rates on Mediterranean buffalo (*B. bubalis*) sperm motility at thawing. *Proc. Intern. Symp. "Mediterranean Animal Germplasm and Future Human Challenges"*, Benevento, Italy, vol. 1 pp. 117-124.
- Dhali, A., R. S. Manik, S. K. Das, S. K. Singla and P. Palta.

- 2000a. Vitrification of Buffalo (*Bubalus bubalis*) oocytes. *Theriogenology*. 1295-1303.
- Dhali, A., R. S. Manik, S. K. Das, S. K. Singla and P. Palta. 2000b. Effect of ethylene glycol concentration and exposure time on post vitrification survival and *in vitro* maturation rate of buffalo oocytes. *Theriogenology*. 53:253. (Abstr).
- Dhaliwal, G. S. and R. D. Sharma. 1990. Serum progesterone profile in buffaloes following two routes of prostaglandin administration. *Indian J. Anim. Sci.* 60:967-968.
- Dhami, A. J. and S. B. Kodagali. 1990. Freezability, enzyme leakage and fertility of buffalo spermatozoa in relation to the quality of semen ejaculates and extenders. *Theriogenology* 34:853-863.
- Dhami, A. J. and K. L. Sahni. 1994. Effects of various cooling (from 30°C to 5°C) equilibration and diluent treatment on freezability, post thaw thermo-resistance, enzyme leakage and fertility of bubaline spermatozoa. *Buffalo J.* 2:147-159.
- Dhami, A. J., V. R. Jain, G. Mohan and K. L. Sahni. 1994. Effect of extenders and additives on freezability, post thaw thermo-resistance and fertility of frozen Murrah buffalo semen under tropical climate. *Buffalo J.* 1:35-45.
- Diaz, J. S., M. Fritsch and J. L. Rodrigues. 1994. A. I. Prefixed in water buffaloes with synchronized estrus using prostaglandin. *Proc. 4th World Buffalo Congress. Sao Paulo, Brazil 27-30 June*, 3:588-590.
- D'Occhio, M. J. D., D. Jillella and B. R. Lindsey. 1999. Factors that influence follicle recruitment, growth and ovulation during ovarian superstimulation in heifers: opportunities to increase ovulation rate and embryo recovery by delaying the exposure of follicles to LH. *Theriogenology* 51:9-35.
- Drost, M., J. M. Wright, W. S. Gipe and A. R. Richter. 1983. Embryo transfer in water buffaloes (*Bubalus bubalis*). *Theriogenology* 20:579 (Abstr).
- Drost, M. 1991. Training manual for embryo transfer in water buffaloes. *FAO Animal Production and Health Paper. No. 84*, pp. 85.
- El Gaffary, M. N., A. E. Abdel Ghafer and L. M. Maral. 1997. Production of buffalo embryos *in vitro*: Applications and limitations. In *International conference on animal, poultry and rabbit production and health. Egyptian International Centre for Agriculture Dokki, Cairo, Egypt 2-4 September (EDICA)*, pp. 113-133.
- El-Amrawi, G. A. 1997. Effect of thawing time and post-thaw temperature of fertility of buffalo spermatozoa frozen in straws. *Proc. 5th World Buffalo Congr., Italy, Vol. I* pp. 850-855.
- Eyestone, W. H. and N. L. First. 1989. Co-culture of early cattle embryos to the blastocyst stage with oviductal tissue or in conditioned media. *J. Reprod. Fert.* 85:715-720.
- Galli, C., G. Crotti, C. Notari, P. Turini, R. Duchi and G. Lazzari. 2001. Embryo production by ovum pick up from live donors. *Theriogenology* 55:1341-1357.
- Galli, C., R. Duchi, G. Croffi and G. Lazzari. 1998. Embryo production by ovum pick up in water buffalo. *Theriogenology* 49:400 Abstr.
- Gandhi, R. S. 1994. Enhanced rates of genetic improvement in Murrah buffaloes through MOET. *Buffalo J.* 3:279-287.
- Gandolfi, F. and R. Moor. 1987. Stimulation of early embryonic development in sheep by co-culture with oviduct epithelial cells. *J. Reprod. Fert.* 81:23-28.
- Gasparrini, B., G. Neglia, R. Di. Palo, G. mpanile and L. Zicarelli. 2000. Effect of cysteamine during *in vitro* maturation on buffalo embryo development. *Theriogenology* 54:1537-1542.
- Gerber, D., T. Arlotto and D. Cooper. 2000a. Comparison between two transvaginal oocyte recovery sessions three days apart in African buffalo (*Synocerus caffer*) 14th International Congress on Animal Reproduction, Sweden, Abstracts Vol. 2:177. Abstr.
- Gerber, D., T. Arlotto and D. Cooper. 2000b. A new disposable needle guide system for transvaginal oocyte recovery in domestic cows and African buffalo. *Theriogenology* 53: 333 Abstr.
- Gordon, I. 1997. *Controlled Reproduction in Buffaloes*. CAB International Wallingford, U.K.
- Gupta, K. A. 1998. Studies on vaginal electrical resistance and plasma progesterone profile in relation to oestrus cycle in buffaloes. *M.V.Sc. Thesis submitted to Rajasthan Agricultural University, Bikaner, India*.
- Gupta, K.A. and G. N. Purohit. 2001. Use of vaginal electrical resistance to predict estrus and ovarian activity, its relationship with plasma progesterone and use for insemination in buffaloes. *Theriogenology* 56:235-245.
- Gupta, P. S. P., S. Nandi, B. M. Ravindranatha and P. V. Sharma. 2001. Isolation of preantral follicles from buffalo ovaries. *Vet. Record*. 148:543-544.
- Haman, A. M., A. A. Salama, M. R. Shalash and I. M. Shihata, 1992. Pharmacological effects of gonadotropic hormones on ovarian activity with special reference to superovulation in buffaloes. *Egyptian J. Vet. Sci.* 29:61-71.
- Hassan, S. G., K. E. L. Battawy, A. A. Menoufy, M. Younis and R. M. Khattab. 2000. Studies on postpartum anestrus in Egyptian buffaloes: Effect of melatonin on ovarian activity 14th International Congress on Animal Reproduction. Stockholm, Abstracts Vol. 1, pp. 185 (Abstr).
- Heap, R. B. and G. C. W. Spencer. 1998. Animal biotechnology : convergence of science, law and ethics. In *Proc. Special Symposium and Plenary Sessions 8th WCAP, Seoul, Korea*, pp. 100-114.
- Heap, W., 1891. Preliminary note on the transplantation and growth of mammalian ova within a uterine foster mother. Cited by S. K. Singla. 1997. In: *In vitro* fertilization, embryo transfer and associated technique in farm animals, EBC, NDRI, Karnal, India, pp. 24.
- Hendolin, M., J. Aalto, K. Kananen Anttila and V. Rainio. 2000. Generation of transgenic male offspring for highly mosaic lactoferrin founder bull using MOET-biopsy-PCR approach. *Proceeding IETS Annual Meeting, Maastricht, The Netherland*.
- Jainudeen, M. R. 1983. Reproduction in the water buffalo: In *Current Therapy in Theriogenology*. W. B. Saunders Company, Philadelphia, pp. 443-444.
- Jainudeen, M. R., 1989. Embryo transfer technology in the buffalo: A review. Paper presented to joint FAO/IAFA/ACTAR Research coordination and planning meeting on Buffalo Productivity, CSIRO, Australia, Feb., 20-24.
- Jainudeen, M. R., Y. Takahashi, M. Nihayah and H. Kanagawa .

- 1993a. *In vitro* maturation and fertilization of swamp buffalo (*Bubalus bubalis*) oocytes. Anim. Reprod. Sci. 31: 205-212.
- Jainudeen, M. R., Y. Takahashi and H. Nehayahm Kangawa. 1993b. *In vitro* maturation and fertilization of swamp buffalo (*Bubalus bubalis*). Theriogenology 37:232 (Abstr).
- Jainudeen, M. R. and E. S. E. Hafez. 1993c. Reproduction in cattle and buffalo. In Reproduction in Farm Animals Edited by ESE Hafez 6th edition Lea and Febriger, Philadelphia.
- Joshi, B. V., S. Rajeshwaran and A. K. Misra. 1992. Effect of FSH priming on superovulatory response in buffalo (*Bubalus bubalis*). Theriogenology 37:232 (Abstr).
- Kamonpatana, M., 1990. Multiovation in buffaloes leading to ET. The world's prospect and achievement at year 1990. Buffalo J., 6:1-10.
- Karaivanov, C. 1986. Comparative studies on the superovulatory effect of PMSG and FSH in water buffalo (*Bubalus bubalis*). Theriogenology 26:51-59.
- Kasiraj, R., A. K. Misra, M. Mutha Rao, R. S. Jaiswal and N. S. Rangareddi. 1993. Successful culmination of pregnancy and live birth following the transfer of frozen thawed buffalo embryos. Theriogenology 39:1187-1192.
- Kasiraj, R., M. M. Rao, N. S. Rangareddy and A. K. Misra. 1992. Superovulatory response in buffaloes following a single subcutaneous or multiple intramuscular FSH administration. Theriogenology 37:234.
- Kaul, G., G. S. Sharma, B. Singh and K. K. Gandhi. 2000. Capacitation and acrosome reaction in buffalo bull spermatozoa assessed by chlortetracycline and *Pisum sativum* agglutinin fluorescence assay. Theriogenology 55:1457-1468.
- Kidson, A., H. Lambrechts, P. Bartels, R. S. Denniston and D. M. Barry. 2000. *In vitro* embryo development in the cape buffalo (*Syncerus caffer*). Age, parity and oocyte quality dependent differences following IVF. Theriogenology 53: 335 Abstr.
- Kitiyanant, Y., J. Saikhum, M. Jaroensuwan, S. Thaiying and K. Pavasuthipaisit. 2000. Nuclear transfer of buffalo fetal fibroblasts and oviductal cells into enucleated bovine oocytes and their subsequent development. Theriogenology. 53:226 (Abstr).
- Kohram, H., G. Mohammadi, N. D. Moradi, M. Sheikhi and M. T. Beigi Nassiri. 2000. Emergence of follicular waves with gonadotropin releasing hormone in buffalo. Theriogenology 53:273 Abstr.
- Kumar, A., V. S. Solanki, S. K. Jindal, V. N. Tripathi and G. C. Jain. 1997. Oocyte retrieval and histological studies of follicular population in buffalo ovaries. Anim. Reprod. Sci. 47:189-195.
- Kumar, D., P. Bissa, P. Goswami and G. N. Purohit. 2002. Effect of epidermal and insulin like growth factor-1 on cumulus expansion, nuclear maturation and fertilization of buffalo oocyte cumulus complexes in serum free media DMEM and Ham's F-10 Compendium 9th International Congress on Biotechnology in Animal Reproduction on 2-4 Dec., Chennai, pp. 224-225 (Abstr.).
- Kumar, P., C. R. Balkrishnan and B. R. Yadav. 1994. Effect of calcium ions on *in vitro* capacitation and acrosome reaction in spermatozoa of buffalo. In Proc. 4th World Buffalo Congress, Sao Paulo, Brazil. Vol. 3, pp. 558-559.
- Kumar, S., Sahani, K.L., Benjamin, B.N. and Mohan, G., 1993. Effects of various levels of yolk on deep freezing and storage of buffalo semen in different diluters without adding glycerol. Buffalo J. 1:79-85.
- Lambrechts, H., E. E. Van Niekerk, W. A. Coetzer, S. W. P. Coleti and G. Vander Horst. 1999. The effect of cryopreservation on the survivability, viability and motility of epididymal African buffalo (*Syncerus caffer*) spermatozoa. Theriogenology 52:1241-1249.
- Lohan, I. S., M. S. Saini, M. L. Kaker, M.L., B. Singh, R. K. Malik and S. S. Grewal. 2000. Follicular changes in post partum buffaloes induced to cyclicity. Theriogenology 52:501 Abstr.
- Lonergan, P., C. Carolan, A. Van Longendonck, I. Donnay, H. Khatri and P. Mermillod. 1996. Role of epidermal growth factor in bovine oocyte maturation and pre implantation embryo development *in vitro*. Biol. Reprod. 54:1420-1429.
- Lu, G. and T. T. Hsu. 1991. *In vitro* fertilization of oocytes in Taiwan water buffalo In Proc. of 3rd World Buffalo Congress, Varma, Bulgaria, Vol. III, pp. 604-609.
- Luizda Silva, A., E. Costa G oncalves, M. P. Schneider B. Breing and A. Krempler. 2002. Construction and analysis of genomic library of the water buffalo (*Bubalus bubalis*). Abstracts XXII World Buiartrics Congress 18-23 August Hannover Germany, pp. 20.
- Madan, M. L. 1984. Superovulation and ova collection. Studies on physiology of buffalo and cattle. 1st National Animal Physiology Workers Conference, NDRI Karnal, India. pp. 26.
- Madan, M. L., S. K. Singla, S. Jaikhani and J. D. Ambrose. 1991. *In vitro* fertilization in buffalo and birth of first ever IVF buffalo calf. Proc. of the 3rd Buffalo Congress. Verma, Bulgaria, 15-19 May, Vol. 7, pp. 11-17.
- Madan, M. L., M. S. Chauhan, S. K. Singla and R. S. Manik. 1994a. Pregnancies established from water buffalo (*Bubalus bubalis*) blastocysts from *in vitro* matured, *in vitro* fertilized oocytes and co-cultured with cumulus and oviductal cells. Theriogenology 42:591-600.
- Madan, M. L., S. K. Singla and R. S. Manik. 1994b. *In vitro* production and transfer of embryos in buffaloes. Theriogenology 41:139-143.
- Mahmoudzadeh, A. R., V. A. Soon, P. Bols, M. J. Ysebaert and A. De., Krizeif. 1996. Optimization of a simple vitrification procedure for bovine embryos produced *in vitro*. Effect of development stage, two step addition of cryoprotectant and semen dilution on embryonic survival, J. Reprod. Fertility. 103:33-39.
- Mamongan, V. G., B. A. Parker, E. B. delos Santos and S. K. Ranjhan. 1989. Proceedings on drought animals in rural development Edited by D Hoffmann et al. ACIAR Proceedings No. 27, Canberra, Australia. pp. 190-194.
- Manik, R. S., M. L. Madan and S. K. Singla. 1992. Ultrasonography in animal production and reproduction. Prod. 12th Int. Congress Anim. Reprod. August 23-27. Hague, Netherlands, pp. 234.
- Markandeya, N. M., D. R. Pargaonkar and S. A. Bakshi. 1993. Studies on vaginal electrical resistance associated with estrus in buffaloes. Buffalo J. 9:157-158.

- Matharoo, J. S. and M. Singh. 1994a. Seasonal variation in superovulatory response and non surgical embryo recovery in buffalo. In Proceedings 4th World Buffalo Congress, Sao Paulo, Brazil, 27-30 June Vol. 3, pp. 652-654.
- Matharoo, J. S. and M. Singh. 1994b. Seasonal variation in the superovulatory response and embryo recovery in the buffalo (*Bubalus bubalis*) Indian J. Anim. Reprod. 18:1-5.
- Mattapallil, M. J. and A. S. Ali. 1998. Parentage assessment of an IVF calf from *Bubalus bubalis* by DNA fingerprinting Gene. 206 (2):209-214.
- McNutt, T. L. and L. A. Johnson. 1996. Flow cytometric sorting of sperm: Influence on fertilization and embryo/fetal development in rabbit. Mol. Reprod. Dev. 43: 261-267.
- Mehmood, A., M. Anwar and M. H. Javad. 1989. Superovulation with PMSG beginning on three different days of the cycle in Nilli Ravi buffaloes. Buffalo J. 5:79-84.
- Misra, A. K., M. C. Yadav and K. T. Motwani. 1988. Successful embryo transfer in a buffalo. Proc. II World Buffalo Congress, New Delhi, India. 1:56.
- Misra, A. K. 1993. Superovulation and embryo transfer in buffaloes : progress, problems and future prospects in India. Buffalo J. 1:13-24.
- Misra, A. K., B. V. Joshi, P. L. Agrawala, R. Kasiraj, S. Sivaiah, N. S. Rangareddi and M. V. Siddiqui. 1992. Cryopreservation of bubaline embryos. Buffalo J. 8:297-303.
- Misra, A. K., R. Kasiraj, M. Mutha Rao, N. S. Ranga Reddy and B. V. Joshi. 1994. Embryo transfer in buffaloes in India. Progress in the last five years. Proc. IVth World Buffalo Congress. 3:501-504.
- Misra, A. K., M. Mutha Rao, R. Kasiraj, N. S. Ranga Reddy and H. C. Pant. 1999. Factors affecting pregnancy rate following non surgical embryo transfer in the buffalo: A retrospective study. Theriogenology 52:1-10.
- Moran, J. B., 1983. Aspects of nitrogen utilization in Asiatic water buffalo and Zebu cattle. J. Agric. Sci. Camb. 100:13-23.
- Muster, E. B., J. Van, Stap, R. A. Hoebe, G. J. Meerman and J. A. Aten. 1999. Difference in sperm head volume as a theoretical basis for sorting X- and Y-bearing spermatozoa : Potentials and limitations. Theriogenology 52:1281-1293.
- Nandi, S., M. S. Chauhan and P. Palta. 1998. Effects of cumulus cells and sperm concentrations on cleavage rate and subsequent embryonic development of buffalo (*Bubalus bubalis*) oocyte matured and fertilized *in vitro*. Theriogenology 50:1251-1262.
- Nguyen, T., N. T. Uoc, L. V. Ty, D. D. Long, D. Chupin, J. P. Renard and B. X. Nguyen. 1997. Effect of estradiol 17- β and hCG supplementation on superovulatory responses and embryo quality in swamp buffalo implanted with norgestomet. Anim. Reprod. Sci. 117:181-187.
- Nilmann, H., 1991. Cryopreservation of ova and embryos from livestock: current status and research needs. Theriogenology. 35:109-123.
- Nityawardana, S., S. Chatchaidy, K. Tantisantikorn, C. Lohacit, P. Chantraprateep and P. Bodhipaksha. 1982. Non surgical embryo transfer in buffaloes. Annual case conference report. Fac. Vet. Sci. Chulalongkorn Univ. Bangkok, Thailand. pp. 23.
- Oba, E., E. J. Fuck, S. D. Bicudo, F. O. Papa and O. M. Ohashi. 1994. Preliminary study on different mediums for deep freezing of buffalo semen. Proc. 4th Int. Buffalo Congr. Sao Paulo Brazil, Vol. 3 pp. 579-581.
- Ocampo, L., M. B. Ocampo, F. P. Aquino, R. De. Vera and I. C. Cruz. 1996. Blastocyst formation of swamp buffalo embryos in co-culture system. Proc. 2nd ABA Congress, Manila, Phillipines. pp. 412-416.
- Ocampo, L.C., F. V. Mamuad, H. Shimizu and M. B. Ocampo. 2001. *In vitro* production of preimplantation buffalo embryos. Buffalo J. 17:145-154.
- Othen, L. S., A. C. Bellem, C. J. Gartley, K. Auckland, W. A. King, R. M. Liptrap and K. I. Goodrowe. 1999. Hormonal control of estrus cyclicity and attempted superovulation in wild bison (*Bison bison athabasca*). Theriogenology 52: 313-323.
- Pal, S. and Dhanda, O.P., 1994. *In vitro* capacitation of buffalo spermatozoa. In Proc. 4th World Buffalo Congress, Sao Paulo, Brazil. Vol. 3:627-629.
- Palasz, A. T. and R. J. Mapletoft. 1996. Cryopreservation of mammalian embryos and oocytes: recent advances. Biotech. Adv. 19:127-149.
- Palta, P. and M. L. Madan. 1984. Hypophyseal response to GnRH in pregnant buffaloes. NDRI Diamond Jubilee Commemorative. Vol. Part III, pp. 334.
- Palta, P. and M. S. Chauhan. 1998. Laboratory production of buffalo (*Bubalus bubalis*) embryos. Reprod. Fert. Dev. 379-391.
- Palta, P., N. Bansal, B. S. Prakash, R. S. Manik and M. L. Madan. 1998. Follicular fluid inhibin concentrations in relation to follicular diameter and estradiol-17 β . Progesterone and testosterone concentrations in individual buffalo ovarian follicles. Indian J. Exp. Biology. 36:768-774.
- Parnpai, R., V. Timsard, M. Kamonpatana, C. Pansin, S. Sophon, A. Lemsakul and C. R. Austin. 1985. Recovery of swamp buffalo embryos using non surgical technique. Buffalo J. 77-82.
- Parnpai, R., K. Tasripoo and M. Kamonpatana. 2000. Comparison of cloning swamp buffalo embryos using foetal fibroblasts and granulosa cells as donor cells. 14th International Congress on Animal Reproduction, Stockholm 2-6 July. Abstracts. Vol. 2, pp. 241.
- Pavasuthipaisit, K., Y. Kitiyanant, C. Thonabulsombat, C. Tocharus, S. Sriurairatna and K. L. White. 1992. *In vitro* maturation and fertilization of swamp buffalo oocytes and their subsequent development. Theriogenology 38:545-555.
- Purohit, G. N., M. Datt and S. S. Sharma. 2000. Embryo recoveries to different superovulatory regimes and live births subsequent to non surgical embryo transfer in Rathi and its crossbred cows. Indian J. Anim. Sci. 70:1064-1066.
- Purohit, G. N. 2001. Effect of serum free media and growth factors on IVM and IVF and intra vaginal culture of buffalo cumulus oocyte complexes. Ph.D. Thesis, Rajasthan Agricultural University, Bikaner, Rajasthan, India.
- Purohit, G. N. and S. S. Sharma. 2002a. *in vitro* maturation of bubaline oocytes in the bovine vagina in serum-free medium. Theriogenology 57:738 (Abstr.).

- Purohit, G. N. and S. S. Sharma. 2002b. *In vitro* maturation of bubaline oocytes in serum-free media in the *in vivo* bovine vagina. *Vet. Rec.* 151 (21):640-641.
- Purohit, G. N. and S. S. Sharma. 2002c. Influence of epidermal growth factor and Insulin like growth factor-I on nuclear maturatin and fertilization of buffalo cumulus cells enclosed oocytes in serum free media and their subsequent development *in vitro*. XXVII Annual Convention and National Symposium on Reproductive Technologies for Augmentation of Fertility in Livestock, Nov. 14-16, IVRI, U.P., India, pp. 45-46 (Abstr.).
- Ramadan, A. A., S. A. Selim. H. M. Harsan and M. A. Wahba. 2000. Immune regulation of ovarian function in buffaloes. *Thriogenology* 55:346-53.
- Ranjhan, S. K. and N. N. Pathak. 1993. Textbook on Buffalo Production. 3rd edition, Vikas Publishing House Pvt. Ltd., New Delhi, India.
- Rao, A. V. and P. Venkatramaiah. 1989. Luteolytic effect of a low dose of cloprostenol monitored by changes in vaginal resistance in subestrus buffaloes. *Anim. Reprod. Sci.* 21:149-152.
- Rao, A. V. N. and M. C. Venkataramulu. 1994. Effect of single varsus double insemination on conception rate in bovines *Indian Vet. J.* 71:1144-1145.
- Rao, K. H. and A. V. N. Rao. 1988. Luteolytic effect of a small dose of cloprostenol administered via intra vulvar route in riverine buffaloes. *Proc. II World Buffalo Congress*, New Delhi, India. Vol. III, pp. 159-161.
- Rao, R. A. 1979. Treatment of subestrus in buffaloes with cloprostenol. *Vet. Rec.* 105, 168.
- Reiger, D., B. Grisart, E. Sample, A. Van Langendonkt, K. J. Betteridge and F. Dessey. 1995. Comparison of the effects of oviductal cell co-culture and oviductal cells conditioned medium on the development and metabolic activity of cattle embryos. *J. Reprod. Fert.* 105:91-98.
- Rorie, R. W. 1999. Effect of timing of Artificial insemination on sex ratio. *Theriogenology* 52:1273-1280.
- Samad, H. A. and A. A. Naserri. 1979. Quantitative study of primordial follicles in buffalo heifer ovaries. *Compendium 13th FAO/SIDA International Course of Animal Reproduction Uppsala, Sweden.*
- Samad, H. A., I. Q. Khan, N. U. Rehman and N. Ahmad. 1998. The recovery, *in vitro* maturation and fertilization of Nili Ravi buffalo follicular oocytes. *Asian-Aust. J. Anim. Sci.* 11:491-497.
- Samad, H. A., N. Akhtar, Alis, N. V. Rahman and Zia, Ur Rahman. 2000. Studies on *in vitro* production of early Nili Ravi buffalo embryos. 14th International Congress of Animal Reproduction, Stockholm, Sweden, Abstracts Vol., 2, p. 207.
- Sansone, G., M. J. F. Natri and A. Fabbrocini. 2000. Storage of buffalo (*Bubalus bubalis*) semen. *Anim. Reprod. Sci.* 62:55-76.
- Shi, D. S., S. F. Yang, S. J. Tan and K. H. Lu. 2000. Effects of enucleation on the parthenogenetic development of bovine and buffalo oocytes. 14th International Congress on Animal Reproduction. Abstracts Vol. 1, p. 29.
- Simonsen, B. T., H. R. Siegismund and P. Arctander. 1998. Population structure of Africn buffalo inferred from mt DNA sequences and microsatellite loci; high variation but low differentiation. *Molecular Ecology.* 7:225-237.
- Singh, B., V. D. Dixit, V. P. Dixit, P. Singh, G. C. George and I. S. Lohan. 2000. Changes in plasma levels of inhibin and FSH in buffaloes superovulated with eCG. *Asian-Aust. J. Anim. Sci.* 13:1205-1209.
- Singh, G., S. M. Totey and G. P. Talwar. 1989. *In vitro* fertilization of oocytes matured *in vitro*. *Theriogenology.* 31:255 (Abstr).
- Singh, R. and A. C. Majumdar. 1992. Chronological changes of buffalo follicular oocyte maturation *in vitro*. *Indian J. Anim. Sci.* 62:205-209.
- Singla, S. K., 1996. Multiplying embryos through nuclear transfer in *Bubalus bubalis*. National Workshop on Application of Tissue Culture in Genetic Studies. March, 17-27, 1996. NDRI Karnal, India. pp. 55-56.
- Singla, S. K. 1997. Nucleus transfer and embryo cloning technique in buffalo gametes. In: *In vitro* fertilization, embryo transfer and associated technique in farm animals. E.B.C., NDRI, Karnal, India. pp. 117-132.
- Singla, S. K., R. S. Manik and M. L. Madan. 1997. Micromanipulation and cloning studies on buffalo oocytes and embryos using nucleus transfer. *Indian J. Exp. Biology.* 35:1273-1283.
- Smith, R. L., K. Lubbe, P. Bartels and R. A. Godke. 1999. Evaluating *in vitro* maturation time and *in vitro* fertilization rate of African buffalo (*Syncerus caffer*) oocytes. *Theriogenology.* 51:294 (Abstr).
- Songsasen, N., V. Yiengvisavakul, B. Buntarach and S. Pharee. 1999. Effect of treatment with recombinant bovine somatotrophin on responses to superovulatory treatment in swamp buffalo (*Bubalus bubalis*). *Theriogenology* 52:377-384.
- Sood, S. K. 1997. PCR Based sexing assay for buffalo embryos. In: *In vitro* fertilization, embryo transfer and associated technique in farm animals. Embryo Biotechnology Centre, NDRI Karnal, India. pp. 86-105.
- Sood, S. K., M. S. Chauhan and O. S. Tomer. 2001. A direct duplex PCR assay for sex determination of Murrah buffalo embryos. *Buffalo. J.* 17:113-124.
- Suzuki, T., S. K. Singla, J. Sujata and M. L. Madan. 1991. Cleavage capability of water buffalo follicular oocytes classified by cumulus cells and fertilized *in vitro*. *J. Vet. Med. Sci.* 53:475-478.
- Suzuki, T., S. K. Singla, J. Sujata and M. L. Madan. 1992. *In vitro* fertilization of water buffalo follicular oocytes and their ability to cleave *in vitro*. *Theriogenology.* 38:1187-1194.
- Tajik, P., S. Goorani Nejad and H. Ghasemzadeh Nava. 2000. *In vitro* maturation of water buffalo oocytes in different concentration of bovine and buffalo follicular fluid. *Theriogenology* 53:470 (Abstr).
- Taneja, M., G. Singh, S. M. Totey and A. Ali. 1995. Follicular dynmics in water buffaloes superovulated in presence or absence of dominant follicle. *Theriogenology* 44:581-597.
- Tavares, L., M. E. O. A. Assumpcao, M. R. B. D. Mella, M. Angelo, P. S. Basurelli, L. M. Queiroz and J. A. Vusinton. 1997. A twice weekly ultrasound guided follicular aspiration in buffalo heifers. *Proc. V. Buffalo Congress.*

- Caserta, Italy. pp. 783-786.
- Techakumphu, M. C., W. Lohachit, W. Tantasuparak, C. Srianan, Intaramongkaol and P. Chantraprateep. 1995. Ovarian response and oocyte recovery in prepubertal swamp buffalo (*Bubalus bubalis*) calves after FSH or PMSG treatment. *Theriogenology* 44:244 (Abstr).
- Thungtanawat, P., D. Abraprasert, S. Keilsommat, E. Aobachit, P. Chantraprateep and P. Bhodipaksha. 1981. Induction of superovulation in buffaloes and non surgical embryo collection. Annual case conference report, Fac. Vet. Sci. Chulalongkorn Univ. Bangkok, Thailand, pp. 16.
- Totey, S. M., G. Singh, M. Taneja, C. H. Pawshe, G. Singh and R. S. Chillar. 1991. Non-surgical embryo transfer in crossbred and non-descript cows. *Indian J. Anim. Reprod.* 12:42-46.
- Totey, S. M., G. Singh, M. Taneja, C. H. Pawshe and G. P. Talwar. 1992. *In vitro* maturation, fertilization and development of follicular oocytes from buffalo (*Bubalus bubalis*). *J. Reprod. Fert.* 95:597-607.
- Totey, S. M., C. H. Pawshe and G. P. Singh. 1993a. Effects of bull and heparin and sperm concentrations on *in vitro* fertilization of buffalo (*Bubalus bubalis*) oocyte matured *in vitro*. *Theriogenology* 39:887-898.
- Totey, S. M., C. H. Pawshe and G. P. Singh. 1993b. *In vitro* maturation and fertilization of buffalo oocytes (*Bubalus bubalis*): Effects of media, hormones and sera. *Theriogenology* 39:1153-1171.
- Totey, S. M., M. Dalvi, K. B. C. Appa Rao, C. H. Pawshe, M. Taneja and R. S. Chillar. 1996. Differential cleavage and development rates and their correlation of cell number and sex ratios in buffalo embryos generated *in vitro*. *Theriogenology* 45:521-523.
- Uoc, N.T., B. X. Nguyen, L. V. Ty, D. Chupin, J. F. Beck and J. P. Renard. 1992. Effect of estradiol supplementation on superovulation in swamp buffalo. *Theriogenology* 38:471-478.
- Vale, W. G. 1997. News on reproductive biotechnology in males. *Proc. 5th World Buffalo Congr. Italy Vol. I* pp. 103-123.
- Venitkul, M. 1989. Progress on embryo transfer in buffalo (Thailand). *Cattle Monthly.* 2:78-81.
- Villa, E. and G. Fabbri. 1993. Results of AI in Italy for improvement schemes in buffalo. *Proc. Int. Symp. on Prospect of Buffalo Production in Mediterranean and the Middle East, Cairo Egypt, Vol. I*, pp. 367-369.
- Vlakhov, K., M. Karaivanov, P. Petrov and D. Kacheva. 1986. Superovulation and the production of embryos in the water buffalo (*Bubalus bubalis*) in Bulgaria. *Vet. Metd. Nauki*, 23:84-88.
- Wall, R. J. 1996. Transgenic livestock-Progress and prospects for future. *Theriogenology* 45:57-68.
- Wilmut, I., A. E. Schnieke, J. McWhir, A. J. Kind and K. H. S. Campbell. 1997. Viable offspring derived from foetal and adult mammalian cells. *Nature.* 385:810-813.
- Windsor, D. P., G. Erans and I. G. White. 1993. Sex predetermination by separation of X- and Y-chromosome bearing sperm: A review. *Reprod. Fert. Dev.* 5:155-171.
- Wright, R. W. Jr. and K. R. Bondioli. 1981. Aspects of *in vitro* fertilization and embryo culture in domestic animals. *J. Anim. Sci.* 5:702-703.
- Xiao Yongzuo. 1989. Production performance of crossbred buffaloes in China. *Proceedings of seminar on buffalo genotypes for small farms in Asia*. Edited by M. K. Vidyadaran et al., Hoffmann et al., ACIAR Proceedings No. 27, Canberra, Australia. pp. 190-194.
- Yadav, B. R., P. K. Katiyar, M. S. Chauhan, M. L. Madan. 1997. Chromosome configuration during *in vitro* maturation of goat, sheep and buffalo oocytes. *Theriogenology* 47:943-951.
- Zicarelli, L., R. Boni, G. Campanile, S. Roviello, B. Gasparrani and R. Di Palo. 1997a. Response to superovulation according to parity, OPU priming, presence/absence of dominant follicle and season. *Prod. V. World Buffalo Congress, Caserta, Italy*, pp. 748-753.
- Zicarelli, L., C. De Filippo, M. Francillo, C. Pacelli and E. Villa. 1997b. Influence of insemination technique and ovulation time on fertility percentage in synchronised buffaloes. *Proc. 5th World Buffalo Congr., Italy, vol. I*, pp. 732-737.